

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

Gram Staining of isolate TW-S-7-1






FIG. 1




Electrophoresis of avian collagen on a 7.5% polyacrylamide gel. α and β denote different collagen configurations.

FIG. 2



Separation of porcine collagen on a 10% polyacrylamide gel. α , β and γ denote different collagen configurations.
Lanes 1, 2 porcine collagen, 20 μ g and 40 μ g; lanes 3, 4 commercial bovine collagen, 40 μ g and 20 μ g


FIG. 3



8% SDS-PAGE

Lane 1:
10 μ l
Lane 2:
20 μ l

FIG. 4



8% SDS-PAGE


Lane 1: 5 μ l

Lane 2: 10 μ l

Lane 3: 20 μ l

Lane 4: 40 μ l

FIG. 5



8% PAGE-SDS

Lane 1 : marker

Lane 2 : 5 μ l

Lane 3 : 10 μ l

FIG. 6